


Claims:

I(We) claim:

- 5 1. An N-substituted piperazine acetic acid active ester compound of the formula:



or a salt thereof, wherein;


LG is the leaving group of an active ester;

X is O or S;

- 10 Y is a straight chain or branched C1-C6 alkyl group or a straight chain or branched C1-C6 alkyl ether group wherein the carbon atoms of the alkyl group or alkyl ether group each independently comprise linked hydrogen, deuterium or fluorine atoms;
- 15 each Z is independently hydrogen, deuterium, fluorine, chlorine, bromine, iodine, an amino acid side chain, a straight chain or branched C1-C6 alkyl group that may optionally contain a substituted or unsubstituted aryl group wherein the carbon atoms of the alkyl and aryl groups each independently comprise linked hydrogen, deuterium or fluorine atoms, a straight chain or branched C1-C6 alkyl ether group that may optionally contain a substituted or unsubstituted aryl group wherein the carbon atoms of the alkyl and aryl groups each independently
- 20 comprise linked hydrogen, deuterium or fluorine atoms or a straight chain or branched C1-C6 alkoxy group that may optionally contain a substituted or unsubstituted aryl group wherein the carbon atoms of the alkyl and aryl groups each independently comprise linked hydrogen, deuterium or fluorine atoms; and
- 25 optionally the N-substituted piperazine acetic acid active ester comprises one or more heavy atom isotopes.
2. The compound of claim 1, wherein the N-substituted piperazine acetic acid active ester is isotopically enriched with one or more heavy atom isotopes.


3. The compound of claim 1, wherein the N-substituted piperazine acetic acid active ester is isotopically enriched with three or more heavy atom isotopes.

- 5 4. The compound of claim 1, wherein LG is:



and wherein X is O or S.


5. The compound of claim 1, wherein LG is:



and wherein X is O or S.

6. The compound of claim 1, wherein LG is N-hydroxysuccinimide.

7. The compound of claim 1, wherein each Z is independently hydrogen, deuterium, fluorine, chlorine, bromine or iodine.
- 5 8. The compound of claim 1, wherein each Z is independently hydrogen, methyl or methoxy.
9. The compound of claim 1, wherein Y is methyl, ethyl, *n*-propyl, isopropyl, *n*-butyl, isobutyl, *sec*-butyl or *tert*-butyl.
- 10 10. The compound of claim 1, wherein X is ^{16}O or ^{18}O .
11. The compound of claim 1, wherein each nitrogen atom of the piperazine ring is independently ^{14}N or ^{15}N .
- 15 12. The compound of claim 1 of the formula:




wherein

- each C* is independently ^{12}C or ^{13}C ;
- LG is the leaving group of an active ester;
- 20 X is O or S;
- Y is a straight chain or branched C1-C6 alkyl group or a straight chain or branched C1-C6 alkyl ether group wherein the carbon atoms of the alkyl group or alkyl ether group each independently comprise linked hydrogen, deuterium or fluorine atoms;
- 25 each Z is independently hydrogen, deuterium, fluorine, chlorine, bromine, iodine, an amino acid side chain, a straight chain or branched C1-C6 alkyl group that may optionally contain a substituted or unsubstituted aryl group wherein the carbon atoms of the alkyl and aryl groups each independently comprise linked hydrogen, deuterium or fluorine atoms, a straight chain or branched C1-C6 alkyl

ether group that may optionally contain a substituted or unsubstituted aryl group wherein the carbon atoms of the alkyl and aryl groups each independently comprise linked hydrogen, deuterium or fluorine atoms or a straight chain or branched C1-C6 alkoxy group that may optionally contain a substituted or unsubstituted aryl group wherein the carbon atoms of the alkyl and aryl groups each independently comprise linked hydrogen, deuterium or fluorine atoms.

13. The compound of claim 2 of the formula:



wherein, LG is the leaving group of an active ester.

14. The compound of claim 13, wherein the compound is a mono-TFA salt, a mono-HCl salt, a bis-TFA salt or a bis-HCl salt.


15. The compound of claim 13, wherein each incorporated heavy atom isotope is present in at least 80 percent isotopic purity.

16. The compound of claim 13, wherein each incorporated heavy atom isotope is present in at least 93 percent or isotopic purity.

17. The compound of claim 13, wherein each incorporated heavy atom isotope is present in at least 96 percent or isotopic purity.


18. The compound of claim 13, wherein LG is N-hydroxysuccinimide.

19. The compound of claim 13, wherein LG is:



and wherein X is O or S.

5 20. The compound of claim 13, wherein LG is:



and wherein X is O or S.

21. The compound of claim 1, wherein the N-substituted piperazine acetic acid active ester is a mono-TFA salt, a mono-HCl salt, a bis-HCl salt or a bis-TFA salt.


22. The compound of claim 2, wherein each incorporated heavy atom isotope is present in at least 80 percent isotopic purity.

23. The compound of claim 2, wherein each incorporated heavy atom isotope is present in at least 93 percent isotopic purity.

5 24. The compound of claim 2, wherein each incorporated heavy atom isotope is present in at least 96 percent isotopic purity.

25. A method comprising:


reacting an N-substituted piperazine acetic acid compound of the formula:



10

or a salt thereof,


with: 1) a compound of the formula:



and, if the piperazine compound is a salt, 2) optionally with a base strong enough to deprotonate the basic nitrogen atoms of the piperazine ring;

15

to thereby form an N-substituted piperazine acetic acid active ester of the formula:



or a salt thereof wherein;

Hal is a fluorine, chlorine, bromine or iodine;

20

LG is the leaving group of an active ester;

X is O or S;

Y is a straight chain or branched C1-C6 alkyl group or a straight chain or branched C1-C6 alkyl ether group wherein the carbon atoms of the alkyl group or alkyl

ether group each independently comprise linked hydrogen, deuterium or fluorine atoms;

each Z is independently hydrogen, deuterium, fluorine, chlorine, bromine, iodine, an amino acid side chain, a straight chain or branched C1-C6 alkyl group that may optionally contain a substituted or unsubstituted aryl group wherein the carbon atoms of the alkyl and aryl groups each independently comprise linked hydrogen, deuterium or fluorine atoms, a straight chain or branched C1-C6 alkyl ether group that may optionally contain a substituted or unsubstituted aryl group wherein the carbon atoms of the alkyl and aryl groups each independently comprise linked hydrogen, deuterium or fluorine atoms or a straight chain or branched C1-C6 alkoxy group that may optionally contain a substituted or unsubstituted aryl group wherein the carbon atoms of the alkyl and aryl groups each independently comprise linked hydrogen, deuterium or fluorine atoms; and optionally the N-substituted piperazine acetic acid moiety comprises one or more heavy atom isotopes; and

optionally treating the N-substituted piperazine acetic acid active ester with an acid.

26. The method of claim 25, wherein the N-substituted piperazine acetic acid active ester is isotopically enriched with one or more heavy atom isotopes.

27. The method of claim 25, wherein the N-substituted piperazine acetic acid active ester is isotopically enriched with three or more heavy atom isotopes.


28. The method of claim 25, wherein the acid is HCl or TFA.

29. The method of claim 26, wherein each incorporated heavy atom isotope is present in at least 80 percent isotopic purity.

30. The method of claim 26, wherein each incorporated heavy atom isotope is present in at least 93 percent isotopic purity.

31. The method of claim 26, wherein each incorporated heavy atom isotope is present in at least 93 percent isotopic purity.


32. The method of claim 25, wherein LG is:



and wherein X is O or S.

5

33. The method of claim 25, wherein LG is:




and wherein X is O or S.

10

34. The method of claim 25, wherein LG is N-hydroxysuccinimide.

35. The method of claim 25, wherein each Z is independently hydrogen, deuterium, fluorine, chlorine, bromine or iodine.

36. The method of claim 25, wherein each Z is independently hydrogen, methyl or methoxy.
37. The method of claim 25, wherein Y is methyl, ethyl, *n*-propyl, isopropyl, *n*-butyl, isobutyl,
 5 *sec*-butyl or *tert*-butyl.
38. The method of claim 25, wherein X is ^{16}O or ^{18}O .
39. The method of claim 25, wherein each nitrogen atom of the piperazine ring is
 10 independently ^{14}N or ^{15}N .
40. The method of claim 25, wherein the compound to be reacted has the formula:




wherein,

- 15 each C* is independently ^{12}C or ^{13}C ;
 LG is the leaving group of an active ester;
 X is O or S;
 Y is a straight chain or branched C1-C6 alkyl group or a straight chain or branched
 C1-C6 alkyl ether group wherein the carbon atoms of the alkyl group or alkyl
 20 ether group each independently comprise linked hydrogen, deuterium or
 fluorine atoms;
 each Z is independently hydrogen, deuterium, fluorine, chlorine, bromine, iodine, an
 amino acid side chain, a straight chain or branched C1-C6 alkyl group that may
 optionally contain a substituted or unsubstituted aryl group wherein the carbon
 25 atoms of the alkyl and aryl groups each independently comprise linked
 hydrogen, deuterium or fluorine atoms, a straight chain or branched C1-C6 alkyl
 ether group that may optionally contain a substituted or unsubstituted aryl
 group wherein the carbon atoms of the alkyl and aryl groups each independently
 comprise linked hydrogen, deuterium or fluorine atoms or a straight chain or

branched C1-C6 alkoxy group that may optionally contain a substituted or unsubstituted aryl group wherein the carbon atoms of the alkyl and aryl groups each independently comprise linked hydrogen, deuterium or fluorine atoms.


- 5 41. The method of claim 23, wherein the product of the reaction is an N-methyl piperazine acetic acid active ester of the formula:



wherein, LG is the leaving group of an active ester.


- 10 42. The method of claim 41, wherein each incorporated heavy atom isotope is present in at least 80 percent isotopic purity.
43. The method of claim 41, wherein each incorporated heavy atom isotope is present in at least 93 percent isotopic purity.
- 15 44. The method of claim 41, wherein each incorporated heavy atom isotope is present in at least 96 percent isotopic purity.

45. The method of claim 41, wherein LG is:



and wherein X is O or S.

5 46. The method of claim 41, wherein LG is:



and wherein X is O or S.

47. The method of claim 41, wherein LG is N-hydroxysuccinimide.

48. The method of claim 41, wherein the N-substituted piperazine acetic acid active ester is a mono-TFA salt, a mono-HCl salt, a bis-HCl salt or a bis-TFA salt.

49. The method of claim 25, wherein the N-substituted piperazine acetic acid active ester is a mono-TFA salt, a mono-HCl salt, a bis-HCl salt or a bis-TFA salt.